

GUAZALIL SL

Reg. No. L 7962

Act No. 36/1947

A soluble concentrate fungicide for post-harvest treatment of citrus for control of Green mould, Blue mould and Sour rot.

ACTIVE INGREDIENTS:

Imazalil as sulphate (imidazole) 134 g/l

Guazatine (guanidine) 200 g/l

Registered by, ICA International Chemicals (Pty) Ltd, Reg. No. 2001/013319/07
P.O. Box 2312, Stellenbosch, 7601.



WARNINGS

- Handle with care.
- Poisonous when swallowed, inhaled or absorbed through skin.
- May irritate skin and cause serious damage to eyes.
- Toxic to fish, other aquatic organisms and wildlife.
- Store in cool place away from food, feed, food utensils and drinking water.
- Keep out of reach of children, animals and uninformed people.
- In case of poisoning call a physician and provide this label and MSDS.

Although this remedy has been extensively tested, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease to the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Do not inhale spray mist.
- Avoid contact with skin and eyes.
- Wear rubber gloves and face shield when handling product.
- Wash with soap and water after use or skin contact. Wash contaminated clothes.
- Do not eat, drink or smoke whilst handling product or before washing hands and face.
- In case of poisoning flush mouth with plenty of water, give lots of water to drink and seek immediate medical assistance. Provide MSDS.
- Note to physician: No specific antidote known. Treat symptomatically. Perform gastric lavage and administer activated charcoal.
- Prevent contamination of food, feed, eating utensils, drinking water, rivers and dams.
- Rinse empty container three times with a volume of water equal to 10% of container. Add the rinsings to the contents of the treatment tank before destroying the container by perforation and burying. Do not use container for other purposes.

RESISTANCE WARNING

For resistance management **Guazalil SL** is a group code 3 and M7 fungicide. Any fungus population may contain individuals naturally resistant to **Guazalil SL** and other group code 3 and M7 fungicides. The resistant individuals can eventually dominate the population if these are used repeatedly. These resistant fungi may not be controlled by **Guazalil SL** or any other group code 3 and M7 fungicide. To delay fungicide resistance:

- Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide group codes.
- For tank mixing or alternation with products in fungicide group code 1, refer to applicable individual product labels.
- Integrate the control methods (chemical, cultural , biological) into disease control programmes.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE

USE ONLY AS DIRECTED

MIXING INSTRUCTIONS:

1. Shake container before use.
2. Agitate the solution during application.
3. To restore the original volume in the dip tank, add required water and then the appropriate amount of **Guazalil SL**.
4. Do not use **Guazalil SL** in waxes.

| CROP & DISEASE | DOSAGE | REMARKS |
|---|--|---|
| CITRUS Post-harvest decay control of Green mould (<i>Penicillium digitatum</i>), Blue mould (<i>Penicillium italicum</i>), and Sour rot (<i>Geotrichum candidum</i>) | Dip treatment: 5 l / 1000 l water (Give a treatment concentration of 500 p.p.m. Imazalil and 1000 p.p.m. Guazatine) | Treat fruit after washing and cleaning. The drenching time is not of critical importance. Water temperature should not exceed 47°C. Do not rinse fruit after treatment. This treatment will also result in good <i>Penicillium</i> sporulation control. |

GENERAL INSTRUCTIONS

1. Handle fruit carefully to minimise infection sites caused by injury.
2. Ensure diseased fruit is promptly removed from the field and packing house and is destroyed.
3. It is recommended that fruit are first washed and treated in a dip tank with **Sporekill** (L7115).
4. Treat fruit promptly after harvest.
5. **Guazalil SL** can be used without wetting agent in water with a pH below 7.
6. Compliance of "directions for use" will ensure that residues do not exceed local Maximum Residue Limits, but the import tolerances of other countries might possibly be exceeded. If the fruit to be treated is intended for export, consult the relevant importer or exporting body regarding the use of this product, Maximum Residue Limits.