

WAARSKUWINGS:

Hanteer versigtig. Giftig deur inname of wanneer deur vel geabsorbeer. Mag oë en vel irriteer. Giftig vir vis, ander akwatiese organismes en wild. Stoor in koel plek weg van voedsel en voer. Hou buite bereik van kinders, oningeligte persone en diere.

Alhoewel hierdie middel omvattend onder 'n groot verskeidenheid toestande getoets is, waarborg die registrasiehouer nie dat dit onder alle toestande doeltreffend sal wees nie, aangesien die werking en effek daarvan beïnvloed kan word deur faktore soos abnormale klimaats- en bergingstoestande; kwaliteit van verdunningswater, verenigbaarheid met ander stowwe wat nie op die etiket aangedui is nie en die voorkoms van weerstand van siekte teen die betrokke middel sowel as die metode, tyd en akkuraatheid van toediening. Verder aanvaar die registrasiehouer nie verantwoordelikheid vir skade aan gewasse, plantegroei, die omgewing of vir enige nadelige effek op mens of vir 'n gebrek aan prestasie van die betrokke middel as gevolg van die versuim deur die gebruiker om die etiketaanwysings na te kom of as gevolg van die ontstaan van toestande wat nie kragtens die registrasie voorsien kon word nie. Raadpleeg die verskaffer in geval van enige onsekerheid.

VOORSORGMATREËLS:

Voorkom inasemend van spuitmis of stof. In geval van toevallige velkontak was met seep en water. Was besoedelde klere. Moenie eet, drink of rook tydens toediening of vermenging nie, of voordat gesig en hande gewas is nie. Voorkom besoedeling van riviere en damme. Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei. Vernietig leë houers deur dit vol gate te maak en plat te slaan. Moet vir geen ander doel gebruik word nie.

BESTANDHEIDSWAARSKUWING:

ImazaCure 750 SG is 'n groepkode 3 swamdoder. Enige populasie van 'n spesifieke swam mag individueel insluit wat 'n natuurlike weerstand teen **ImazaCure 750 SG** of enige ander groepkode 3 swamdoder het. Indien hierdie swamdoders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die swampopulasie oorheers. Hierdie weerstandbiedende swamme sal waarskynlik nie deur **ImazaCure 750 SG** of enige ander groepkode 3 swamdoder beheer word nie. Om weerstand teen swamdoders te vertraag:

1. Vermoë die eksklusiewe herhaaldelike gebruik van swamdoders met dieselfde groepkode. Wissel af met of gebruik tenkingsels van produkte in verskillende swamdoder groepkodes.
2. Raadpleeg toepaslike individuele produk-etiketke wanneer daar afgewissel word, of tenkingsels gemaak word, met produkte in swamdoder groepkodes M en U.
3. Integreer ander beheermaatreëls (chemies, verbouing en biologies) in swamdoder programme.

Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

ImazaCure 750 SG word aanbeveel vir die beheer van na-oesbederf van sitrusvrugte veroorsaak deur *Penicillium digitatum* (groenskimmel) en *Penicillium italicum* (blou-skimmel).

ImazaCure 750 SG kan gebruik word sonder benattingsmiddels in water met pH laer as 7.

TOEDIENING EN DOSIS: Moet nie ImazaCure 750 SG in wakse gebruik nie.**A. Aanwending in waterbad.**

Meng 670g **ImazaCure 750 SG** met 1000 liter water om 'n behandelingskonsentrasie van 500 d.p.m. Imazalil te verkry. Die temperatuur van die water moet nie 47°C oorskry nie. Sorg dat die swamdoder goed opgelos is. Die indompelingstyd is nie van kritiese belang nie.

B. Nie-herwinbare aanwending deur sproei- of berseltoediening.

Voeg 34g **ImazaCure 750 SG** by 25 liter water en roer totdat swamdoder opgelos is. Dit gee 'n behandelingskonsentrasie van 1000 d.p.m. Imazalil. Die 25 liter **ImazaCure 750 SG** oplossing moet gebruik word om hoogstens 17000 kg vrugte te behandel.

Moet nie vrugte afspoel na behandeling nie. Hierdie behandelings sal ook goeie sporulasie beheer tot gevolg hê.

IMAZACURE

750 SG

Reg. No. : L 7926

Wet / Act 36 / 1947

'N WATEROPLOSBAAR KORREL SWAMDODER VIR NA-OESBEHANDELING VAN SITRUS**A WATER SOLUBLE GRANULE FUNGICIDE FOR POST-HARVEST TREATMENT OF CITRUS.**

FRAC FUNGICIDE GROUP 3 FRAC SWAMDODER GROEP

AKTIEWE BESTANDDEEL / ACTIVE INGREDIENT

Imasalil as sulfaat / Imazalil as sulphate (imidasool/imidazole) _____ 750g/kg

Batch No:

Date of
Manufacture:.....

Nett Contents

670g

Registrasiehouer / Registration holder :

ICA International Chemicals (PTY) Ltd • Reg. No. 2001/013319/07 • P.O. Box 2312, Stellenbosch, 7601

UN No. 2588

CAUTION
VERSIGTIG**WARNINGS:**

Handle with care. Poisonous when swallowed or absorbed through the skin. May irritate eyes and skin. Toxic to fish, other aquatic organisms and wildlife. Store in a cool place away from food and feed. Keep out of reach of children, uninformed persons and animals.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

Avoid inhalation of spray mist and dust. In case of skin contact, wash thoroughly with soap and water. Wash contaminated clothing. Do not eat, drink or smoke whilst applying or preparing the spray mix or before washing face and hands. Prevent contamination of rivers and dams. Prevent contamination of food, feed, drinking water and eating utensils. Destroy empty container by perforation and flattening and do not use for any other purpose.

RESISTANCE WARNING:

For resistance management **ImazaCure 750 SG** is a group 3 fungicide. Any fungus population may contain individuals naturally resistant to **ImazaCure 750 SG** and other group 3 code fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly. The resistant fungi may not be controlled by **ImazaCure 750 SG** or any other group 3 fungicides. To delay fungicide resistance:

1. Avoid exclusive repeated use of fungicides from the same fungicide group code. Alternate or tank mix with products from different fungicide codes.
2. For tank mixing or alternation with products in fungicide group code M or U, refer to applicable product labels.
3. Integrate control methods (chemical, cultural and biological) into disease control programs.

For specific information on resistance management contact the registration holder of this product.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

ImazaCure 750 SG is recommended for control of post harvest decays of citrus fruits caused by *Penicillium digitatum* (green mould) and *Penicillium italicum* (blue mould).

ImazaCure 750 SG can be used without wetting agent in water with a pH below 7.

APPLICATION AND DOSAGE RATE: Do not use **ImazaCure 750 SG** in waxes.**A. Water bath treatment.**

Add 670g **ImazaCure 750 SG** to 1000 liters of water to obtain a treatment concentration of 500 p.p.m. Imazalil. See that fungicide is well dissolved. Water temperature should not exceed 47°C. The drenching time is not of critical importance.

B. Non-recovery water spray-on or brush-on.

Add 34g **ImazaCure 750 SG** to 25 liters water and mix until dissolved. This gives a treatment concentrate of 1000 p.p.m. Imazalil. The 25 liters **ImazaCure 750 SG** solution must be used to treat not more than 17000 kg of fruit.

Do not rinse fruit after treatment. These treatments will also result in good sporulation control.